

量子物理学・ナノサイエンス第 400 回セミナー

**Proton-induced knockout ($p,2N$) reactions on stable nuclei
as a tool to determine spectroscopic factors**

講師： 若狭 智嗣 教授
九州大学 大学院理学研究院

日程： 11 月 13 日（水）14:00 -

場所： 本館 1 階 M102 講義室

概 要

Proton-induced knockout reactions provide a direct mean of studying the single particle or cluster structures of target nuclei. In this talk, firstly we review the experimental and theoretical progress in this field. The spectroscopic factors extracted using ($p,2N$) data at intermediate energies are consisted with those determined using ($e,e'p$) data, typically within a 15% deviation, when the geometrical parameters are the same as those employed in the ($e,e'p$) analysis. However, when the goal is to use the ($p,2N$) reactions as a spectroscopic tool, it is preferable to determine the geometrical parameters in a self-consistent manner. Thus we further performed a calculation using wave functions generated in a relativistic Hartree model. The spectroscopic factors deduced from this self-consistent calculation and from the previous standard calculation using a global optical potential agree with the relevant ($e,e'p$) results mostly within 15% for light nuclei. This result is encouraging for extending to similar studies of unstable nuclei, for which the properties are not well-known and the parameters for DWIA calculations cannot be pre-determined.

※「物理学特別講義（発展）第二」を履修する学生は本セミナーも聴講すること。

連絡教員 関口 仁子（内線 2298）